

Ф. Л. Быков¹, В. А. Гордин^{2,1}

¹ФБГУ «Гидрометцентр России»
²НИУ «Высшая школа экономики»

**Комплексный коэффициент турбулентного обмена по данным
зондирования с высоким разрешением атмосферы Земли
(модификация модели Аккерблома — Экмана)**

17 декабря 2021 г


The traditional **Akerblom – Ekman model** describes the dynamics of wind speed in the boundary layer (BL) of the atmosphere or an ocean on a rotating planet:

$$\begin{cases} \frac{d}{dz} \left[k(z) \frac{du}{dz} \right] = -l (v - v_g), \\ \frac{d}{dz} \left[k(z) \frac{dv}{dz} \right] = l (u - u_g), \end{cases} \quad (1)$$

where $u(z)$, $v(z)$ are the required horizontal wind components, u_g, v_g are the geostrophic wind on the BL upper boundary. The vertical variable $z \in [0, H_{\max}]$ is the height above the Earth's surface, H_{\max} is the thickness of the boundary layer, $l = \sin \varphi \times 1.45842 \times 10^{-4} / s$ is the Coriolis parameter, φ is the geographic latitude, $k(z) > 0$ is the coefficient of the turbulent exchange. **The system (1) is singular iff $k(z_0) = 0$ for some $z_0 \in (0, H_{\max})$.**

If $k(z) = const$ then the wind rotation angle in BL is equal to 45° .

Observed wind rotation angle – histogram



Conditional probability distributions of the wind rotation angle

Eq. (1) is invariant with respect to the group of rotations around the vertical axis $\mathbf{SO}(2)$. The group $\mathbf{SO}(2)$ commutes only with skew-symmetric operators. Therefore, we will also consider the more general system:

$$\begin{cases} \frac{d}{dz} \left[\gamma(z) \sin(\varphi) \frac{dv}{dz} + k(z) \frac{du}{dz} \right] = -l (v - v_g), \\ \frac{d}{dz} \left[k(z) \frac{dv}{dz} - \gamma(z) \sin(\varphi) \frac{du}{dz} \right] = l (u - u_g), \end{cases} \quad (2)$$

where the second coefficient of turbulent exchange $\gamma(z)$ plays the role of the regularizator for (1): when $k(z)=0$, the system (2) does not become degenerate. The cofactor $\sin(\varphi)$ was added into Eq. (2) to adjust the results of our numerical experiments with data from South hemisphere.

The complex form

Let us rewrite (2) in a complex form: $w = u + \mathbf{i}v$, $w_g = u_g + \mathbf{i}v_g$, and $\kappa = k - \mathbf{i}\gamma \sin(\varphi)$:

$$\frac{d}{dz} \left[\kappa(z) \frac{dw}{dz} \right] = \mathbf{i}l (w - w_g). \quad (3)$$

If $\kappa(z) = \text{const}$ then the wind rotation angle is equal $\arg \sqrt{\frac{\mathbf{i}}{\kappa}} = \frac{1}{2} \text{atan} \frac{k}{\gamma \sin \varphi}$.

The observable wind rotation angle $10 - 20^\circ$ corresponds to the values of the ratio

$$\gamma \sin \varphi / k \approx 1.2 - 2.7$$

The quadratic programming problem (QPP)

To reduce the order of differentiation, we integrate Eq. 2 with respect to z :

$$\kappa(z) \frac{dw}{dz} = -\psi + c \quad (4)$$

where $c \in \mathbb{C}$ are constant of integration, a function $\psi(z)$ satisfy the following equations:

$$\frac{d\psi}{dz} = il(w_g - w), \quad \int_0^H \psi(z) dz = 0.$$

We will search $\kappa(z)$ as a solution of QPP. This QPP minimizes the mean relative residual of (4) over N vertical profiles:

$$L(\kappa(z), \{c_j\}) = \frac{1}{N} \sum_{j=1}^N \frac{1}{W_j} \int_0^{H_j} \left| \kappa(z) \frac{dw_j}{dz} + \psi_j(z) - c_j \right|^2 dz \rightarrow \min_{\kappa(z), \{c_j\}}, \quad (5)$$

$W_j = \int_0^{H_j} |\psi(z)|^2 dz$ for the normalization. Then $\min_{c_j} L(0, c_j) = 1$.


Let $0 < \Lambda < 1$ be the minimum of the functional (5). The value $100\% (1 - \Lambda)$ is interpreted as the average coefficient of determination. It is presented in Table 2.

Dataset

We use the dataset from **26142** profiles, which satisfy the following conditions:

1. The measurement accuracy for wind speed is $0.1m/s$ (not 1 knot).
2. The mean vertical resolution is good (more than 25 points in the layer 0-1000 m).
- 3. The boundary layer thickness $H_j > 100m$.**
4. The variability of the wind in the boundary layer is greater than $2.5m/s$.
5. The height of the first measurement in the BUFR profile is no more $5m$.

Geographical location of the aerological stations



Geographical location of 111 stations, from which the radiosonde data were assimilated. Crosses mark **28** “intensive” stations, with a large number (more than 400) of the profiles

Boundary layer's thickness

We use a standard definition of the boundary layer's thickness H_j as the minimal positive root of the following equation:

$$\Theta_j(H_j) = \Theta_{V,j}(0), \quad (5)$$

where Θ is a potential temperature and Θ_v is a potential virtual temperature.

The dataset of BUFR profiles during the period from Apr. 4, 2018 to Nov. 29, 2019

| Subsample name | Addition condition | Profiles | Boundary layer thickness H_j |
|----------------|---|----------|--------------------------------|
| Full | None | 26142 | 671±516m |
| Deep | $H_j > 1000m$ | 8462 | 1592±454m |
| Thin | $H_j < 500m$ | 12051 | 270±115m |
| Stable | $Ri(z) > 0.3$ for any $z \in [0; H_j]$ | 2622 | 201±105m |
| Unstable | $Ri(z_0) < 0.2$ for some $z_0 \in [0; H_j]$ | 22584 | 742±517m |

The Richardson number


The Richardson number Ri is the dimensionless function of the height z :

$$Ri(z) = \frac{g}{\Theta} \frac{\frac{\partial \Theta}{\partial z}}{\left(\frac{\partial u}{\partial z}\right)^2 + \left(\frac{\partial v}{\partial z}\right)^2}. \quad (6)$$


The values $Ri > Ri_c = 0.25$ correspond to stable stratifications, $Ri < Ri_c$ correspond to unstable ones, and $Ri < 0$ correspond to strictly unstable stratifications of an atmospheric column (a temperature inversion layer exists in the column).

Results

a) the real part $k(S_1)$,



b) the imaginary part $\gamma(S_1)$.



The optimal (for different subsamples) coefficient of turbulent exchange κ depending on the relative height $S_1 = z/H$:


a) the real part $k(S_1)$;

b) the imaginary part $\gamma(S_1)$.


We consider also normalized

$$\text{parameters: } \kappa = \frac{\tilde{\kappa} \cdot H_j}{1000m}$$

a) the real part $\tilde{k}(S_1)$,





b) the imaginary part $\tilde{\gamma}(S_1)$.



The optimal normalized coefficients of turbulent exchange $\tilde{k}(S_1)$ depending on the relative height $S_1 = z / H_j$, were determined separately for 28 “intensive” stations.

The wind rotation of the model angle for the optimal $\kappa = H \tilde{\kappa}(z/H)$







Optimal coefficient of turbulent exchange \tilde{k} (full subsample): the real part $\tilde{k}(S_2)$ (graphs 1, 3); the imaginary part $\tilde{\gamma}(S_2)$ (graphs 2, 4) depending on the wind shear module

$$S_2 = shear(z) = \sqrt{(u(z) - u_g)^2 + (v(z) - v_g)^2}$$

($m \cdot s^{-1}$):

for stable subsample (graphs 1, 2),

for unstable subsample (graphs 3,4)



Optimal coefficients of turbulent exchange \tilde{k} (Full subsample) depending on the $S_3 = \arctan(Ri)$, where Ri – Richardson number.

Black vertical line shown the critical Richardson number $Ri = Ri_c = 0.25$. For $Ri > 0.5$ we have the ratio $\gamma / k > 20$

| Optimized coefficients | | \mathcal{K} | $k > 0, \gamma = 0$ | $\tilde{\mathcal{K}}$ | $\tilde{k} > 0, \gamma = 0$ | Ratio |
|------------------------|------------------------------|---|---------------------|-----------------------|-----------------------------|---|
| Subsample | Atmospheric parameter(s) S | The mean coefficient of determination $100\%(1 - \Lambda)$ | | | | $\frac{1 - \Lambda(\tilde{\mathcal{K}} \in \mathbb{C})}{1 - \Lambda(\tilde{\mathcal{K}} \in \mathbb{R})}$ |
| Full | Relative height S_1 | 38,5% | 11,7% | 48,3% | 13,8% | 3,5 |
| Deep | | 34,6% | 7,7% | 35,0% | 8,0% | 4,5 |
| Thin | | 62,3% | 17,4% | 72,6% | 20,9% | 3,3 |
| Stable | | 65,9% | 10,2% | 77,5% | 11,6% | 7,1 |
| Unstable | | 38,9% | 12,3% | 48,2% | 14,0% | 3,5 |
| Full | Wind shear modulus S_2 | 37,6% | 12,2% | 46,3% | 15,1% | 3,1 |
| Deep | | 29,2% | 8,3% | 29,8% | 8,6% | 3,4 |
| Thin | | 59,9% | 18,2% | 67,7% | 22,4% | 3,0 |
| Stable | | 59,7% | 11,6% | 67,8% | 13,5% | 5,5 |
| Unstable | | 39,1% | 12,6% | 46,7% | 15,3% | 3,0 |
| Full | Richardson number Ri | 24,4% | 11,4% | 34,1% | 12,6% | 2,7 |
| Deep | | 29,2% | 7,4% | 29,9% | 7,5% | 4,1 |
| Thin | | 36,0% | 16,8% | 43,0% | 19,8% | 2,1 |
| Stable | | 33,0% | 9,7% | 39,4% | 10,3% | 4,0 |
| Unstable | | 28,6% | 11,8% | 35,3% | 12,8% | 2,7 |

| | | | | | | |
|----------|--|-------|-------|--------------|-------|------------|
| Full | Relative height S_1 and wind shear modulus S_2 | 41,2% | 12,7% | 53,4% | 15,4% | 3,4 |
| Deep | | 35,2% | 8,7% | 35,7% | 9,1% | 4,0 |
| Thin | | 65,2% | 18,9% | 76,7% | 22,9% | 3,3 |
| Stable | | 66,3% | 11,7% | 78,1% | 13,5% | 6,3 |
| Unstable | | 42,0% | 13,1% | 53,4% | 15,7% | 3,4 |
| Full | Relative height S_1 and Richardson number Ri | 39,5% | 12,4% | 50,3% | 13,8% | 3,6 |
| Deep | | 34,7% | 7,9% | 35,1% | 8,1% | 4,5 |
| Thin | | 63,6% | 18,4% | 74,7% | 21,4% | 3,4 |
| Stable | | 65,9% | 10,2% | 77,5% | 11,6% | 7,1 |
| Unstable | | 40,6% | 12,9% | 50,1% | 14,1% | 3,6 |

Comparison of the BUFR profiles and model's solutions

Let us represent the coefficient of the turbulent exchange \mathcal{K} in the form $\mathcal{K} = H \tilde{\kappa}(S_1)$. Then we can find the solution $\hat{w}_j(z, \mathcal{K}, w_0)$ of Eq. 2 with the Dirichlet boundary conditions $w(H) = w_g$, $w(0) = w_0$ and estimate the mean error of the profile reconstruction:


$$ABS_{speed}(S_1, \mathcal{K}, w_0) = \frac{1}{N} \sum_{j=1}^N \left| \left| \hat{w}_j(S_1 H_j, \mathcal{K}, w_0) \right| - \left| w_j(S_1) \right| \right|,$$

$$ABS_{direction}(S_1, \mathcal{K}, w_0) = \frac{1}{\tilde{N}} \sum_{j=1}^{\tilde{N}} \left| \arg \hat{w}_j(S_1 H_j, \mathcal{K}, w_0) - \arg w_j(S_1) \right|,$$


Here we exclude from the formulas for $ABS_{direction}$ the terms with small velocities $|\hat{w}_j|$ or $|w_j| \leq 2 \text{ m/s}$, when the determination of the wind's direction is not clear. The limit of the sums in these formulas is smaller: $\tilde{N} \approx 0.69N$.

The profile reconstruction error

for wind speed modulus



for wind speed angle



Conclusion

1. The original theory of Akerblom – Ekman, predicted 45° wind rotation in the boundary layer. We observed the rotation angle is an average of three times smaller.
2. We include the coefficient γ in the model, the consistence with BUFR data increase up to 7 times for stable stratification and up to 3.5 for unstable. The coefficient γ can be interpreted as a coefficient in the imaginary part of coefficient κ ;
3. We compare the universal coefficient κ , both on unique parameter: relative height $S_1 = z/H$, or on the wind shear S_2 , or on the Richardson number Ri . The relative height is preferable

<http://method.meteorf.ru/ansambl/ansambl.html>

Thank you for attention

This article was prepared within the framework of the Academic Fund Program at the National Research University Higher School of Economics (HSE) in 2020 - 2021 (grant № 20-04-021) and by the Russian Academic Excellence Project 5-100